ELECTRON-POSITRONIUM SCATTERING AND FESHBACH RESONANCES

A S Ghosh

Department of Theoretical Physics, Indian Association for the Cultivation of Science, Jadavpur, Kolkata-700 032, India

This progress report details the scattering of electrons by positronium (Ps) atoms and Feshbach resonances up to n=4 threshold due to doubly excited auto detaching states of positronium negative ion (Ps⁻) carried out by our group [1,5].

It is of interest to study the spectra of Ps^- and compare it with that of H^- . Present dissertation provides the details of close coupling approximation (CCA) model employed by us. This article also discusses the other elaborate methods already employed to investigate the system [2-4]. Comparisons amongst the predictions by different methods constitute a section in this article.

References

- [1] A. Basu and A. S. Ghosh, Europhys. Lett. 60, 46 (2002).
- [2] Y. K. Ho, Phys. Lett. 120A, 348 (1984); Y. K. Ho and A. K. Bhatia, Phys. Rev. A 45, 6268 (1992); I. Ivanov and Y. K. Ho, Phys. Rev. A 60, 1015 (1999) and the references cited therein.
- [3] J. Botero and C. H. Green, Phys. Rev. Lett., 56, 1366 (1986).
- [4] S. J. Ward, J. W. Humberston and M. R. C. McDowell, J. Phys. B 20, 127 (1987).
- [5] A. Basu and A. S. Ghosh (Communicated).